

REDUCED TRANSITION PROBABILITIES TO THE FIRST 2^+ STATE IN $^{52,54,56}\text{Ti}$ *

D.-C. Dinca^{a,b)}, R. V. F. Janssens^{c)}, A. Gade^{b)}, D. Bazin^{b)}, R. Broda^{d)}, C. M. Campbell^{a,b)},
M. P. Carpenter^{c)}, P. Chowdhury^{c,e)}, J. M. Cook^{a,b)}, A. Deacon^{f)}, B. Fornal^{d)}, S. J. Freeman^{f)},
T. Glasmacher^{b)}, F. G. Kondev^{c)}, J.-L. Lecouey^{b)}, S. N. Liddick^{a,b)}, P. F. Mantica^{b)},
W. F. Mueller^{b)}, H. Olliver^{a,b)}, K. Starosta^{a,b)}, J. R. Terry^{a,b)}, B. A. Tomlin^{a,b)}, and K. Yoneda^{b)}

^{a)}*Department of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA*

^{b)}*National Superconducting Cyclotron Laboratory, Michigan State University, East Lansing, MI 48824, USA*

^{c)}*Physics Division, Argonne National Laboratory, Argonne, IL 60439, USA*

^{d)}*Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Krakow, Poland*

^{e)}*Department of Physics, University of Massachusetts, Lowell, MA 01854, USA*

^{f)}*Schuster Laboratory, University of Manchester, Manchester M13 9PL, UK*

The neutron-rich Ti nuclei have been the subject of much interest recently because of experimental evidence for a sub-shell closure at $N = 32$ in neutron-rich nuclei just above ^{48}Ca [1,2] and the development of a new effective interaction GXPF1 [3] predicting a shell gap at $N = 34$. Interestingly, and perhaps surprisingly, the latter calculations do not appear to be substantiated by a recent beta-decay study of the level structure of ^{56}Ti [4].

In order to gain further insight in these nuclei, an experiment was undertaken at the NSCL where the titanium isotopes with $A = 52, 54$, and 56 were studied using intermediate-energy Coulomb excitation in inverse kinematics. $^{52,54,56}\text{Ti}$ fragments were produced via fragmentation of $130 \text{ MeV/A } ^{76}\text{Ge}^{30+}$ primary beam at about 9 pnA average intensity on a ^9Be target. Following separation of the desired fragments in the A1900 separator [5], the beam was directed onto a ^{197}Au target. The Coulomb excitation process was tagged by requiring a coincidence between scattered Ti nuclei identified in the S800 magnetic spectrograph [6] and gamma rays detected with the Segmented Germanium Array (SeGA) [7].

The measured $B(E2; 0^+ \rightarrow 2^+)$ rates will be presented and compared with results of shell model calculations using a number of modern effective interactions.

*This work was supported by the National Science Foundation under Grant Nos. PHY-00110253, PHY-9875122, and PHY-0244453, the U.S. Department of Energy, Office of Nuclear Physics under contract W31-109-ENG-38, by the UK EPSRC, and by the Polish Scientific Committee.

- [1] J. I. Prisciandaro *et al.*, Phys. Lett. **B510**, 17 (2001).
- [2] R. V. F. Janssens *et al.*, Phys. Lett. **B546**, 55 (2002).
- [3] M. Honma *et al.*, Phys. Rev. C **65**, 061301 (2002).
- [4] S. Liddick *et al.*, Phys. Rev. Lett. **92**, 072502 (2004).
- [5] D. J. Morrissey *et al.*, Nucl. Instrum. Methods **B203**, 90 (2003).
- [6] D. Bazin *et al.*, Nucl. Instrum. Methods **B204**, 629 (2003).
- [7] W. F. Mueller *et al.*, Nucl. Instrum. Methods **A466**, 492 (2001).